

1 **Patterns of locomotor regulation during the pole vault approach phase**

2 **Laurie Needham, Timothy A. Exell, Ian N. Bezodis, and Gareth Irwin**

3 **Abstract:**

4 A successful approach phase is key to achieving high performances in the pole vault. The aim of this
5 study was to explore the nature of locomotor control patterns during the pole vault approach phase.
6 Fourteen well-trained athletes performed ten jumps which were recorded using 2D video sampling at
7 200 Hz and analysed. Key kinematics were reconstructed from camera data using a modified 2D-DLT.
8 Patterns of regulation were determined from the standard deviation of footfall locations during the
9 approach phase. These patterns were found to be highly individual but structural differences between
10 those who did and those who did not regulate were identified. Regulation of locomotion was associated
11 with an ability to produce functionally adaptable movement patterns and the consistent achievement of
12 desired performance outcomes. Coaches should include training exercises that require intentional use
13 of regulation to aid athletes in achieving the flexibility to adapt to changing constraints during the
14 approach phase. Athletes should be considered on an individual basis in order to effectively, efficiently
15 and safely improve performance.

16 **Keywords:** Pole vault, approach phase, regulation, adaptability.

17 **Introduction:**

18 Pole vaulting requires athletes to clear a high horizontal cross bar using a flexible vaulting pole. In order
19 to achieve the correct take-off characteristics and maximise the potential to be successful the athlete
20 must satisfy a number of demands during the approach phase. These include concurrently achieving a
21 high horizontal velocity, coordinating the lowering of the pole into the plant box and consistently
22 achieving an accurate take-off position. Various studies have examined different aspects of the pole
23 vault from kinematics (Hay, 1994; Angulo-Kinzler et al., 1994), energetics (Schade, Arampatzis &
24 Brüggemann, 2000; 2004; 2006), and simulation (Hubbard, 1980; Ekevad & Lundberg, 1995; Liu,
25 Nguang & Zhang, 2011) perspectives. Previous research has established that greater peak heights are
26 associated with high horizontal velocities during the approach phase (Greig & Yeadon, 1997;
27 Adamczewski & Perl, 1997; Frere et al., 2010). Frere et al. (2009) concluded that pole carriage caused

28 decreases in running velocity (6.6%) as a result of significantly reduced step lengths in novice athletes,
29 but these findings were from an unconstrained run with no requirement to achieve a desired take-off
30 location or perform the rest of the jump.

31 A reconceptualisation of pole vault performance can be derived from the constraints lead
32 approach (McGinnis & Newell, 1982) which considers the interaction of the athlete, task and
33 environment, based on the Dynamical Systems Theory (DST) (Newell, 1986). Unique to pole vault is
34 the task constraint, created by the need to carry and coordinate the lowering of a vaulting pole and the
35 spatio-temporal constraint created by the necessity to take-off in a specific location (plant box) with the
36 absence of a visual and physical target (e.g. take-off board in long jump and triple jump (Lee, Lisham
37 & Thompson, 1982; Hay & Koh, 1988).

38 The need for the athlete to achieve a precise and consistent take-off location is essential for
39 success. This consistency at take-off can be considered to correspond to the concept of low end-point
40 variability of footfall location, which is considered to be a key performance factor within pole vault
41 coaching literature (Richardson, 2012) as well as for wider gait-regulated disciplines such as long and
42 triple jump (Hay & Koh, 1988). Consistent performance outcomes can be achieved by different patterns
43 of coordination (Bernstein, 1967) and as such, movement pattern variability can be considered
44 functional if it permits the performer the flexibility to adapt to changing constraints during goal-directed
45 actions (Barris, Farrow, & Davids, 2014). The concept of degeneracy provides the theoretical framework
46 to explain functional movement variability and provides athletes with robustness against perturbations
47 (Whitacre & Bender, 2010; Davids et al., 2013; Seifert et al., 2013). Movement patterns can be
48 continuously adapted in a functional way to allow skilled consistent performance outcomes rather than
49 attempting to utilise rigid, stereotyped movement patterns (Barris et al., 2014). Evidence from gait-
50 regulated tasks such as triple-jump (Wilson et al., 2008) demonstrates that individuals are capable of
51 finding different ways to achieve the same performance outcome, even under similar task and
52 environmental constraints. In gait-regulated tasks such as the pole vault approach phase, it has been
53 proposed that performers make adjustments through visual control mechanisms (Lee et al., 1982; Hay,
54 1988; Glize & Laurent, 1997; Bradshaw, 2004) where by the athlete uses perceptual reference points
55 close to the target to control locomotion. This visual information provides a continuous regulation
56 process based on a perception-action coupling (Montagne, Cornus, Glize, Quaine, & Laurent, 2000).

57 Locomotor control mechanisms have been explored extensively within gait-regulated tasks such as

Formatted: Not Highlight

58 long jumping, gymnastics vaulting and walking tasks, and appear to be present across populations,
59 regardless of the athlete's level of skill (Bradshaw & Aisbett, 2006), age (Berg et al., 1994, Panteli et
60 al., 2014), or familiarity with the task (Scott et al., 1997). Typically these control mechanisms have been
61 studied using spatio-temporal variables such as changes in step length and footfall location variability
62 (Lee, Lisham & Thompson, 1982; Hay, 1988) with additional insight being provided by the assessment
63 of the relationship between the adjustments in step length required and adjustments produced to
64 successfully complete the task (Montagne et al., 2000).

65 In the context of pole vaulting, little is known about the approach phase which is more complex
66 in nature than previously studied tasks (e.g. walking, long jump, gymnastics vault etc.) due to additional
67 constraints such as pole carriage, discussed above, and a higher risk of serious injury should the task
68 not be completed correctly. Some evidence (Hay, 1988) exists to support the notion that elite male pole
69 vaulters utilise similar control strategies to other gait regulated tasks but further research is required to
70 assess and understand the strategies of elite and developing skill levels. The aim of this study was to
71 explore the nature of locomotor control patterns during the pole vault approach phase. The purpose of
72 gaining this information was to inform coaches when prescribing approach phase training exercises. It
73 was hypothesised that athletes would present individual patterns of locomotion regulation during the
74 pole vault approach phase.

75

76

77 **Methods:**

78

79 *Participants*

80 Ethical approval was granted by the University's Research Ethics Committee and all participants
81 provided written informed consent. Eleven male (mean \pm SD age: 21 \pm 4 years, height: 1.85 \pm 0.07 m,
82 mass: 76.7 \pm 12.7 kg) and three female athletes (mean \pm SD age: 17 \pm 3 years, height: 1.63 \pm 0.02 m,
83 mass: 60.9 \pm 6.25 kg) were recruited. Performance level was assessed against the current senior world
84 record. Male personal bests ranged between 70% and 90% of the world record while female personal
85 best ranged between 65% and 80% of the world record.

86 *Experimental set-up*

87 Data collections were conducted during a single session at an indoor athletics centre. Kinematic data
88 were collected using four HDV cameras (Type HVR – Z5E; Sony, Japan) placed at a perpendicular
89 angle, 25 m from the approach runway (Figure 1). A sample rate of 200 Hz was selected with a shutter
90 speed of 1/425 s and an open iris. Calibration of the performance area was achieved using a single
91 object of known distances placed sequentially along the centre of the runway to create a 40 m x 3 m
92 plane. Additional recordings were made with a second object consisting of markers of known distances
93 in order to test accuracy and precision of reconstruction.

94

95 ***** FIGURE 1 NEAR HERE *****

96

97 Anthropometric data were collected before participants conducted a self-selected warm-up similar to
98 that normally used during their training sessions. Each participant was required to perform ten jumps
99 over an elastic training bar set between 95-98% of their personal best from a full approach run of self-
100 selected distance. Bar height was determined following discussions with national level coaches. This
101 height range was selected to encourage athletes to perform a regular jump without invoking
102 performance changes that might be associated with attempting to perform jumps at maximal or
103 substantially submaximal heights. Successful jumps (where the athlete attempted to complete a full
104 jump over the bar) were assessed qualitatively by an experienced national level pole vault coach who
105 was present at all data collections. Any trial that was qualitatively deemed to be unsuccessful was
106 discarded. Participants were instructed to allow for full recovery between trials. The number of attempts
107 required to complete the requisite number of jumps was recorded for each athlete. This data was used
108 to determine success rate.

109 *Data Analysis*

110 Camera images were imported to MATLAB (V2013b; The Mathworks Inc. Natick, USA) where an open
111 source digitisation toolbox (Hedrick, 2008) was used to locate the position of desired landmarks. These
112 landmarks included the vertex, C7, hip, shoulder, elbow, wrist, knee, ankle, MTP joint centres and
113 proximal and distal end of the pole. A modified 2D-Direct Linear Transformation (DLT) (Woltring &
114 Huiskes, 1990) was used and a ninth parameter was added to account for the non-linearity of the lens

115 in accordance with Walton (1981). Total body centre of mass (CoM) locations in the vertical (z) and
116 horizontal (y) axes were calculated using de Leva's (1996) model. CoM location of the foot segment
117 was calculated using Winter's (2009) model with an additional mass, determined by weighting each
118 participants shoe, added to account for each individual's footwear mass (Bezodis, 2008). Additionally,
119 pole mass and CoM locations were ascertained using a balance test.

120 For each participant, spatio-temporal characteristics including step velocity (SV), step length
121 (SL) and step frequency (SF) were calculated in accordance with Bezodis et al., (2008). Instances of
122 touch-down and toe-off were identified in order to calculate the duration of ground contact time (GCT)
123 and flight time (FT). Between-trial variability of the toe-to-plant box distance were assessed via the
124 standard deviation of each footfall location in the y-direction (SD_{ff}).

125 Participants were grouped post-hoc as either regulators or non-regulators utilising the regulation
126 definitions of Hay (1988) and Berg et al. (1994). Examples of each pattern are provided in figure 2.
127 These definitions were as follows:

- 128 - Ascending/Descending Pattern (A/D) – An overall increase in the SD_{ff} proceeded by a marked
129 and systematic decrease in SD_{ff} .
- 130 - Ascending Only (AO) – Only, a systematic increase in SD_{ff} is observed.
- 131 - Random Fluctuations (RF) – Small, random-like fluctuations are present in SD_{ff} throughout the
132 approach phase.

133 Based upon these definitions participants were grouped, post-hoc as either regulators or non-
134 regulators. Step numbers are denoted so that 'final' represents the final ground contact, 'penultimate'
135 represents the step immediately preceding the final step, '-3' represents the step preceding the
136 penultimate step... and so on.

137

138 ***** FIGURE 2 NEAR HERE *****

139

140

141 In accordance with previous gait regulation research (Hay, 1988; Montagne et al., 2000; Renshaw
142 & Davids, 2004) SD_{if} for each step, the distribution of adjustments for the final six steps and an intra-
143 step analysis of adjustment required and adjustments produced for the final six steps were calculated.
144 SD_{if} profiles for each step allow for consistency of footfall placement to be mapped across the entire
145 approach phase. Due to the differing approach lengths utilised by participants (12-18 steps) data
146 presented in Figure 3 were time normalised to 101 data points in order to clearly present each
147 individual's SD_{if} pattern. 0% represents the first footfall location of the approach phase i.e. at the end of
148 the first step and 100% represents the end of the approach phase i.e. the take-off step.

Commented [B1]: I think this clarifies the point further

Commented [ET2]: the end of the take off step (just to clarify in relation to lan's previous comment)?

Formatted: Not Highlight

149 Intra-step analysis was conducted by assessing the relationship between the magnitudes of step
150 adjustments required and produced. Adjustment required (Adj_R) were calculated as the difference
151 between the mean footfall location across all trials and the actual footfall location for a given step.
152 Adjustments produced (Adj_P) were calculated as the difference between the mean step length across
153 trials and the actual subsequent step length (Montagne et al., 2000). Linear regression analyses were
154 utilised in order to assess the extent to which performers were capable of producing the required
155 amount of adjustment for each step of the run-up. A Shapiro-Wilk test confirmed that data were normally
156 distributed.

157 In order to explore the underlying structure of variables discussed above for each group, a principle
158 components analysis (PCA) was implemented. Input variables were selected based upon the
159 underlying theory (Hair et al., 2010) utilising variables that describe locomotor regulation during the
160 approach phase. Eight variables were loaded into the PCA input matrix (CoM Velocity, SL, SF, GCT,
161 FT, SD_{if} , Adj_R , Adj_P). Sampling adequacy was confirmed using a Kaiser-Meyer-Olkin test. For each
162 group, data were processed for a PCA using a custom written script in MATLAB (V2016a; The
163 Mathworks Inc. Natick, USA). The number of principle components required to explain 95% of the
164 variance in the data were computed using a Scree test criterion. For each of these identified principle
165 components (PC), a set of component coefficients were also produced. Component coefficients
166 represent the correlation coefficients between the variables and the principles components. Component
167 loadings exceeding ± 0.4 were considered to indicate significant loading (Hemphill, 2003) and any
168 variable which was similarly correlated to multiple components was considered to cross-load, and was
169 therefore discarded from the analysis.

170

171 **Results:**

172 SD_# patterns that were identified to match the A/D pattern (n = 8) were deemed to show evidence of
173 regulation while patterns matching either the R/F (n = 3) or A/O (n = 3) pattern were deemed to not
174 show evidence of regulation based upon this measure. Example SD_# patterns for each regulation
175 definition are shown in figure 2. For the regulation group, 94% of jumps were deemed to be successful
176 while for the non-regulation group, 54% of the jumps were deemed to be successful.

177

178 ***** FIGURE 3 NEAR HERE *****

179

180 For the regulation group mean take-off location accuracy was 0.10 m ± 0.04 m with a maximum
181 SD_# during the approach of 0.15 m ± 0.05 m, while for the non-regulation group, mean take-off location
182 accuracy was 0.09 m ± 0.05 m with a maximum SD_# during the approach of 0.09 m ± 0.05 m. The step
183 for the onset of regulation for the regulation group was between step -5 and -2 while no such step could
184 be identified for the non-regulation group.

185 Intra-step regression analysis described the linear relationship between the amount of Adj_R and
186 the amount of Adj_P. In the regulation group intra-step analysis revealed statistically significant
187 correlation coefficients ($p < 0.05$) between Adj_R and Adj_P at the penultimate and final steps (Figure 4,
188 left). No correlation coefficients were found to be significant in the non-regulation group ($p > 0.05$) at any
189 step (Figure 4, right).

190 ***** FIGURE 4 NEAR HERE *****

191

192 Results of the PCA analysis showed that at least 95% of the variance was accounted for in six
193 and five principle components for the regulation group and non-regulation group respectively. The first
194 principle component accounted for 38% of the variance for the regulation group and 39% of the variance
195 in the non-regulation group.

196 ***** TABLE 1 NEAR HERE *****

197 For the regulation group (Table 1), PC1 and PC3 were most heavily loaded with variables which
198 represent regulation of locomotion (i.e. SD_{ff} and SL on PC1 and Adj_P and Adj_R on PC3). CoM Velocity
199 was found to cross-load between PCs and was discarded. In contrast for the non-regulation group
200 (Table 1), CoM Velocity loaded heavily on PC1. SD_{ff} and Adj_P were cross loaded between PCs.

201 **Discussion and Implications:**

202 Based on the underlying mechanics of the pole vault approach phase and applying the paradigm of
203 Dynamical Systems Theory (DST) this study aimed to explore the nature of locomotor control patterns
204 during the pole vault approach phase. The purpose was to add to the knowledge of regulation of
205 locomotion during complex skills and to inform coaches who prescribe approach phase training
206 exercises.

207 Pole vaulters in this study demonstrated three distinct patterns of SD_{ff} . The majority of pole
208 vaulters in this sample ($n = 8$) presented an A/D pattern while A/O ($n = 3$) and R/F ($n = 3$) patterns were
209 less common. These findings align with previous research in similar gait regulated tasks such as long
210 jumping where the A/D pattern was most common (Hay & Koh, 1988). The A/D pattern was remarkably
211 similar to that observed in previous long jump studies (Lee et al., 1982; Hay & Koh; Scott et al., 1997;
212 Panteli et al., 2014) in terms of the presence of an ascending/descending pattern and the onset point
213 of regulation. This suggests that the majority of pole vaulters did regulate locomotion to achieve a
214 desired take-off location.

215 Regulation patterns do not appear to be associated with skill level here given that the top two
216 performers in this sample presented different patterns. Further to this, performers who demonstrated
217 an R/F pattern presented very low levels of variability throughout the approach phase, demonstrating
218 that high performance levels can be achieved through the use of differing regulation strategies. The R/F
219 regulation strategy is the closest to a stereotyped movement pattern i.e. an approach run with the
220 absence of variability (Richardson, 2013). However, this strategy may lack robustness as these
221 participants do not demonstrate an ability to make functional adjustments during the approach phase,
222 which may be required to ensure success through take-off position consistency. Movement system
223 robustness or the ability to functionally adapt to perturbations in the task are commonly associated with
224 expert behaviour (Seifert et al., 2013). Expert performance has been associated with stable movement
225 patterns that are not stereotyped and rigid but flexible and adaptable, since neurobiological systems

226 can exploit inherent degeneracy (Edelman & Gally, 2001). These concepts are further supported when
227 success rates are considered, see results section. Those who showed evidence of adaptability, i.e.
228 were able to produce a stable movement pattern when needed or a flexible movement pattern when
229 needed (Seifert et al., 2013), achieved a 94% success rate (A/D pattern - regulation group). In contrast,
230 those who showed evidence of a rigidly stable and inflexible movement pattern (A/O or RF pattern -
231 non-regulation group) achieved a 54% success rate. On this basis, the post hoc grouping utilised in this
232 study seem justified. It should be noted that all trials presented in this study were successful ones which
233 may in part explain the similarities in take-off location accuracy between groups.

234 Correlations analysis between Adj_R and Adj_P revealed significant relationships for the
235 penultimate and final steps in the regulation group only. Given that the non-regulation group did not
236 show evidence of regulating or adjusting gait it is unsurprising that no significant correlations were
237 observed. Adjustments produced by the regulation group occurred later during the pole vault approach
238 phase, than during the long jump approach phase (Montagne et al., 2000; Panteli et al., 2014) where a
239 significant correlation was noted at every step after the onset of regulation (approximately six steps
240 from take-off). This later onset of regulation for pole vaulters may be attributed to the reduced
241 accumulation of variance in footfall location (0.15 m) when compared to long jumpers (0.23 m for elite
242 performers (Hay, 1988); 0.29 m for junior performers (Berg *et al.*, 1994)). Lower variability in footfall
243 locations would therefore reduce the demand for regulation. When the pole vault approach phase is
244 considered in the context of a perception-action couple (Glize & Laurent, 1997; Montagne et al., 2000),
245 perceptual information that signifies the need to produce adjustments would be expected to arrive later
246 in the approach phase when magnitudes of variability are lower.

247 The influence of pole carriage upon regulation of gait remains unclear. Where the pole vaulter
248 experiences greater constraints due to the demands of coordinating the lowering of the pole, the
249 flexibility to adapt to local conditions may be limited. Additionally, the high risk of injury associated with
250 not achieving the correct take-off location cannot be ignored (Rebella et al., 2008; Boden et al., 2012).
251 While an inability to adapt and produce adequate adjustments during a long jump approach phase may
252 lead to a discounted jump, failure to produce adequate adjustments during the pole vault approach
253 phase can result in serious injury (Rebella et al., 2008; Boden et al., 2012).

254 In this sample, individual response patterns were present within both groups. Each individual
255 produced a unique set of results in order to satisfy their own intrinsic dynamics (Turvey, 1990). In order

256 to investigate potential driving principles governing the behaviour of the movement system an
257 exploratory PCA was utilised. Structural differences in the data between the regulation group and non-
258 regulation group were identified. For the regulation group, the first three principle components were
259 heavily loaded with variables which describe regulation of gait and velocity, two of the key task demands
260 of the approach phase. In contrast, for the non-regulation group, only velocity based variables loaded
261 onto PCs (Table 1). Two unique data structures were identified, one where the movement system is
262 governed by a combination of velocity and regulatory based variables (regulation group) and one which
263 is governed only by velocity based variables. Structural differences between the two groups were also
264 noted as six PCs accounted for over 95% of the variance in the regulation group data while five PCs
265 were required for the non- regulation group. Increased complexity has been linked to the prevention of
266 the system becoming too stable and thus preventing the emergence of functional movement solutions
267 (Davids et al., 2003). These findings advocate the need for future research to conduct a detailed
268 analysis of the coordinative structures that emerge during the pole vault approach phase under
269 interacting constraints (Seifert et al., 2014). Further, while the influence pole carriage may have an effect upon the
270 findings of this study, it is beyond the scope of this research to understand what this influence may be.
271 Further research, assessing the influence of pole carriage experimentally is therefore required.

272 The results illustrate a clear inability by some performers (non-regulation group) to achieve
273 consistent performance outcomes, in terms of success rates, and explore reasons why these individuals
274 cannot satisfy the regulatory task demands of the pole vault approach phase. By linking the application
275 of biomechanics, motor control and training theory (Dick, 2007), these findings can provide coaches
276 with meaningful information relating to the performer's approach phase performance and facilitate the
277 development of athlete-specific training drills.

278 Practical solutions can be derived from a performer's approach phase data which develop the ability to
279 functionally interact with key constraints (i.e. the task and environment) (Davids et al., 2013). In the
280 pursuit of expert performance, degenerate behaviours (Edelman & Gally, 2001) can be explored to
281 widen the bandwidth of variability that performers can work within while still achieving consistent
282 performance outcomes. When implementing training drills that introduce locomotor regulation and
283 promote functional variability during the approach phase, practitioners should manipulate key task
284 constraints, including perception-action constraints (Davids et al., 2013), that facilitate the emergence
285 of flexible and adaptable movement patterns. For example, for those identified as regulatory athletes,

286 perturbing the approach phase by adjusting the starting position may prove useful. In order to still
287 achieve the desired take-off location the athlete would be required to regulate their approach by differing
288 amounts thus challenging their regulatory ability. In contrast, for athletes identified as non-regulatory,
289 introducing additional perceptual information, such as a clear take-off mark on the runway, might assist
290 in the development of regulatory abilities.

291 **Conclusion:**

292 Pole vaulters in this study demonstrated three distinct patterns of SD_{fr} . Locomotor regulation occurred
293 predominantly during the penultimate and final steps. Patterns of regulation were highly individual but
294 structural differences between those who did and those who did not regulate were identified.
295 Regulation of locomotion was associated with an ability to produce functionally adaptable movement
296 patterns and the consistent achievement of desired performance outcomes. These key findings can
297 be linked to the application of training theory to allow coaching practitioners to prescribe informed
298 interventions in the pursuit of performance enhancement. Athletes should be considered on an
299 individual basis in order to effectively, efficiently and safely improve performance. Future work should
300 consider the robustness of these patterns under changing task constraints.

301 **References:**

- 302
- 303 Adamczewski, H., & Perl, B. (1997). Run-up velocities of female and male pole vaulting and some
304 technical aspects of women's pole vault. *New studies in athletics*, 12, 63-76.
- 305 Angulo-Kinzler, R. M., Kinzler, S. B., Balius, X., Turro, C., Caubet, J. M., Escoda, J., & Prat, J. A. (1994).
306 Biomechanical analysis of the pole vault event. *Journal of Applied Biomechanics*, 10(2), 147-165.
- 307 Barris, S., Farrow, D., & Davids, K. (2014). Increasing functional variability in the preparatory phase of
308 the takeoff improves elite springboard diving performance. *Research Quarterly for Exercise and*
309 *Sport*, 85(1), 97-106.
- 310 Berg, W. P., Wade, M. G., & Greer, N. L. (1994). Visual regulation of gait in bipedal locomotion:
311 Revisiting Lee, Lishman, and Thomson (1982). *Journal of Experimental Psychology: Human Perception*
312 *and Performance*, 20(4), 854.
- 313 Bernstein, N. A. (1967). The co-ordination and regulation of movements. Pergamon Press.

314 Bezodis, I. N., Kerwin, D. G., & Salo, A. I. (2008). Lower-limb mechanics during the support phase of
315 maximum-velocity sprint running. *Medicine and science in sports and exercise*, 40(4), 707.

316 Boden, B. P., Boden, M. G., Peter, R. G., Mueller, F. O., & Johnson, J. E. (2012). Catastrophic Injuries
317 in Pole Vaulters A Prospective 9-Year Follow-up Study. *The American journal of sports medicine*, 40(7),
318 1488-1494.

319 Bradshaw, E. J., & Aisbett, B. (2006). Visual guidance during competition performance and run-through
320 training in long jumping. *Sports Biomechanics*, 5(1), 1-14.

321 Bradshaw, E. (2004). Gymnastics: Target-directed running in gymnastics: a preliminary exploration of
322 vaulting. *Sports Biomechanics*, 3(1), 125-144.

323 Davids, K., Glazier, P., Araújo, D., & Bartlett, R. (2003). Movement systems as dynamical
324 systems. *Sports medicine*, 33(4), 245-260.

325 Davids, K., Araújo, D., Vilar, L., Renshaw, I., & Pinder, R. (2013). An ecological dynamics approach to
326 skill acquisition: Implications for development of talent in sport. *Talent Development & Excellence*, 5(1),
327 21-34.

328 De Leva, P. (1996). Adjustments to Zatsiorsky-Seluyanov's segment inertia parameters. *Journal of*
329 *biomechanics*, 29(9), 1223-1230.

330 Dick, F. W. (2007). *Sports training principles*. A. & C. Black.

331 Edelman, G. M., & Gally, J. A. (2001). Degeneracy and complexity in biological systems. *Proceedings*
332 *of the National Academy of Sciences*, 98(24), 13763-13768.

333 Ekevad, M., & Lundberg, B. (1995). Simulation of 'smart' pole vaulting. *Journal of Biomechanics*, 28(9),
334 1079-1090.

335 Frere, J., Chollet, D., & Tourny-Chollet, C. (2009). Assessment of the influence of pole carriage on sprint
336 kinematics: A case study of novice athletes. *International Journal of Sports Science and*
337 *Engineering*, 3(1), 3-10.

338 Frere, J., L'hermette, M., Slawinski, J., & Tourny-Chollet, C. (2010). Mechanics of pole vaulting: a
339 review. *Sports biomechanics*, 9(2), 123-138.

340 Glize, D., & Laurent, M. (1997). Controlling locomotion during the acceleration phase in sprinting and
341 long jumping. *Journal of sports sciences*, 15(2), 181-189.

342 Greig, M. & Yeadon, M. (1997) *Biomechanics of the Pole Vault*. The Sports council.

343 Hair, J. F., Anderson, R. E., Babin, B. J., & Black, W. C. (2010). *Multivariate data analysis: A global
344 perspective* (Vol. 7). Upper Saddle River, NJ: Pearson.

345 Hay, J. G., & Koh, T. J. (1988). Evaluating the approach in the horizontal jumps. *International Journal
346 of Sport Biomechanics*, 4(4), 372-392.

347 Hay, J. G. (1988). Approach strategies in the long jump. *International Journal of Sport
348 Biomechanics*, 4(2), 114-129.

349 Hay, J. (1978). *The biomechanics of sports techniques*. Prentice-Hall.

350 Hedrick, T. L. (2008). Software techniques for two-and three-dimensional kinematic measurements of
351 biological and biomimetic systems. *Bioinspiration & biomimetics*, 3(3), 034001.

352 Hemphill, J. F. (2003). Interpreting the magnitudes of correlation coefficients.

353 Hubbard, M. (1980). Dynamics of the pole vault. *Journal of Biomechanics*, 13(11), 965-976.

354 Lee, D. N., Lishman, J. R., & Thomson, J. A. (1982). Regulation of gait in long jumping. *Journal of
355 Experimental Psychology: Human perception and performance*, 8(3), 448.

356 Liu, G., Nguang, S. K., & Zhang, Y. (2011). Pole vault performance for anthropometric variability via a
357 dynamical optimal control model. *Journal of Biomechanics*, 44(3), 436-441.

358 McGinnis, P. M., & Newell, K. M. (1982). Topological dynamics: A framework for describing movement
359 and its constraints. *Human Movement Science*, 1(4), 289-305.

360 Montagne, G., Cornus, S., Glize, D., Quaine, F., & Laurent, M. (2000). A perception-action coupling
361 type of control in long jumping. *Journal of motor behavior*, 32(1), 37-43.

362 Needham, L., Bezodis, I., Exell, T., Simpson, S., & Irwin, G. (2016, November). LOCOMOTOR
363 CONTROL PATTERNS DURING THE POLE VAULT APPROACH PHASE. In *ISBS-Conference
364 Proceedings Archive* (Vol. 34, No. 1).

365 Newell, K. M. (1986). Constraints on the development of coordination. *Motor development in children:*
366 *Aspects of coordination and control*, 34, 341-360.

367 Panteli, F. N., Theodorou, A., Piliandis, T., & Smirniotou, A. (2014). Locomotor control in the long jump
368 approach run in young novice athletes. *Journal of sports sciences*, 32(2), 149-156.

369 Rebella, G. S., Edwards, J. O., Greene, J. J., Husen, M. T., & Brousseau, D. C. (2008). A prospective
370 study of injury patterns in high school pole vaulters. *The American journal of sports medicine*, 36(5),
371 913-920.

372 Renshaw, I., & Davids, K. (2004). Nested task constraints shape continuous perception–action coupling
373 control during human locomotor pointing. *Neuroscience Letters*, 369(2), 93-98.

374 Richardson, A. (2013). PV Technical Model: Approach Run [Online]. www.ucoach.com: British
375 Athletics. [2013].

376 Schade, F., Arampatzis, A., & Brüggemann, G. P. (2000). Influence of different approaches for
377 calculating the athlete's mechanical energy on energetic parameters in the pole vault. *Journal of*
378 *Biomechanics*, 33(10), 1263-1268.

379 Schade, F., Arampatzis, A., Brüggemann, G. P., & Komi, P. V. (2004). Comparison of the men's and
380 the women's pole vault at the 2000 Sydney Olympic Games. *Journal of Sports Sciences*, 22(9), 835-
381 842.

382 Schade, F., Arampatzis, A., & Brüggemann, G. P. (2006). Reproducibility of energy parameters in the
383 pole vault. *Journal of biomechanics*, 39(8), 1464-1471.

384 Scott, M. A., Li, F. X., & Davids, K. (1997). Expertise and the regulation of gait in the approach phase
385 of the long jump. *Journal of sports sciences*, 15 (6), 597-605.

386 Seifert, L., Button, C., & Davids, K. (2013). Key properties of expert movement systems in sport. *Sports*
387 *Medicine*, 43(3), 167-178.

388 Seifert, L., Komar, J., Barbosa, T., Toussaint, H., Millet, G., & Davids, K. (2014). Coordination pattern
389 variability provides functional adaptations to constraints in swimming performance. *Sports*
390 *Medicine*, 44(10), 1333-1345.

391 Turvey, M. T. (1990). Coordination. *American psychologist*, 45(8), 938. Whitacre & Bender, 2010.

392 Walton, J. S. (1981) Close-range cine-photogrammetry: a generalised technique for quantifying gross
393 human move- structure. Unpublished Ph.D thesis, College of Health, Physical Education, and
394 Recreation, Pennsylvania State University.

395

396 Wilson, C., Simpson, S. E., Van Emmerik, R. E., & Hamill, J. (2008). Coordination variability and skill
397 development in expert triple jumpers. *Sports biomechanics*, 7(1), 2-9.

398 Winter, D. A. (2009). *Biomechanics and motor control of human movement*. John Wiley & Sons.

399 Woltring, H., & Huiskes, H. (1990). In, *Biomechanics of human movement: applications in rehabilitation,*
400 *sports and ergonomics / Ed. N. Berme, A. Cappozzo. - Worthington, Ohio, 108-127.*

401

402 Figure 1. Schematic diagram showing camera positions relative to the runway. Calibration locations are
403 defined by the crosses, black lines indicate each camera's field of view. (Not to scale).

404

405 Figure 2. Example SD_{if} profiles for each of the regulation types as defined by Hay (1988) and Berg et
406 al. (1994) (adapted from Needham et al., 2016). Solid line, A/D pattern. Dashed line, A/O pattern.
407 Dotted line, R/F pattern.

408

409 Figure 3. Mean SD_{if} profiles for regulation group (left) and non-regulation group (right) athletes with
410 individual profiles provided in gray. Regulation group athletes presented an A/D pattern (left – solid
411 lines) while non-regulation group presented either R/F (right – dashed line) or A/O patterns (right –
412 dashed-dot line).

413

414 Figure 4. The relationship (R^2) between the amount of SL adjustment required and the amount of step
415 SL adjustment produced for each group (left, regulation group & right, non- regulation group). *
416 Indicates significant correlations ($p < 0.05$).

417

418

419

